

UN	목 표	2. End hunger, achieve food security and improved nutrition and promote sustainable agriculture
	세 부 목 표	2.2 By 2030, end all forms of malnutrition, including achieving, by 2025, the internationally agreed targets on stunting and wasting in children under 5 years of age, and address the nutritional needs of adolescent girls, pregnant and lactating women and older persons
	Indicator	2.2.1 Prevalence of stunting (height for age <-2 standard deviation from the median of the World Health Organization (WHO) Child Growth Standards) among children under 5 years of age

I. Global indicator

<Type 2>

Indicator	Prevalence of stunting (height for age <-2 standard deviation from the median of the World Health Organization (WHO) Child Growth Standards) among children under 5 years of age
Definition	Prevalence of stunting (height-for-age < -2 standard deviation from the median of the World Health Organization (WHO) Child Growth Standards) among children under 5 years of age.

II. Data description

[Data] Prevalence of stunting among children under 5 years of age

Calculation method	Prevalence of stunting among children = $\frac{\text{No. of children whose height-for-age is less than } <-2 \text{ standard deviation from the median}}{\text{Population under 5 years of age}} \times 100$
Unit	Percentage(%)
Data sources	Compiled based on the UNICEF global database (CRING).
Calendar	<ul style="list-style-type: none"> ■ Time series: 2000-2019(Data for Korea included in 2003, 2010) ■ Data release: Annually
Data compilers	UNICEF(United Nations Children's Fund), WHO(World Health Organization), WB(World Bank)
Global indicator link	<ul style="list-style-type: none"> ■ Metadata: https://unstats.un.org/sdgs/metadata/files/Metadata-02-02-01.pdf ■ Data: https://unstats.un.org/sdgs/indicators/database/