U N	Goal	3. Ensure healthy lives and promote well-being for all at all ages
	Target	3.9 By 2030, substantially reduce the number of deaths and illnesses from hazardous chemicals and air, water and soil pollution and contamination
	Indicator	3.9.3 Mortality rate attributed to unintentional poisoning

I. National indicator

⟨Type 1⟩

Indicator	Mortality rate attributed to unintentional poisoning
Definition	Deaths per 100,000 population attributed to unintentional poisoning and exposure to noxious substances(X40-X49) * Classification under the seventh Korean Standard Classification of Diseases(KCD-7)
Calculation method	$$\text{Mortality rate attributed to unintentional poisoning} = \frac{\text{Deaths attributed to unintentional poisoning and exposure to noxious substances}}{\text{Mid-year resident population}} \times 100{,}000$$
Unit	Deaths per 100,000 population

II. National indicator's source

Data sources	▌Source: *Cause of Death Statistics* ▌Collection method: Compiling data from death reports filed with city, district, and municipal offices, baby and fetal death reports from crematories, and data from the 「*Complementary Survey on Causes of Death*」 of medical institutions
Calendar	▌Frequency: Annually (Jan 1 - Dec 31 of the reference year) ▌Data release: September in the following year
Organizations	Vital Statistics Division(Tel. 042-481-2252), Statistics Korea
Related International Agency	WHO

III. Comparison with UN SDG indicator

① Indicator		② Definition		③ Data value	
Same	Different	**Same**	Different	Same	**Different**

Note	▌The WHO collects country-specific death data and applies correction if under-reporting is suspected. The scope of deaths attributed to unintentional poisoning includes the ICD codes X40, X43-44, and X46-49
Global indicator link	▌Metadata: https://unstats.un.org/sdgs/metadata/files/Metadata-03-09-03.pdf ▌Data: https://unstats.un.org/sdgs/indicators/database/

SUSTAINABLE DEVELOPMENT GOALS